
CONTENTS

Preface	v
I. The One-Dimensional Hubbard Model	1
[1] E. H. Lieb and F. Y. Wu: <i>Absence of Mott transition in an exact solution of the short-range one-band model in one dimension</i> <i>Phys. Rev. Lett.</i> 20 (1968) 1445.	9
[2] A. A. Ovchinnikov: <i>Excitation spectrum of the one-dimensional Hubbard model</i> <i>Sov. Phys. JETP</i> 30 (1970) 1160.	13
[3] C. F. Coll: <i>Excitation spectrum in the one-dimensional Hubbard model</i> <i>Phys. Rev. B</i> 9 (1974) 2150.	17
[4] F. Woynarovich: <i>Excitations with complex wavenumbers in a Hubbard chain: I. States with one pair of complex wavenumbers</i> <i>J. Phys. C</i> 15 (1982) 85.	26
[5] F. Woynarovich: <i>Excitations with complex wavenumbers in a Hubbard chain: II. States with several pairs of complex wavenumbers</i> <i>J. Phys. C</i> 15 (1982) 97.	38
[6] F. Woynarovich: <i>Spin excitations in a Hubbard chain</i> <i>J. Phys. C</i> 16 (1983) 5293.	51
[7] F. Woynarovich: <i>Low-energy excited states in a Hubbard chain with on-site attraction</i> <i>J. Phys. C</i> 16 (1983) 6593.	63
[8] A. Klümper, A. Schadschneider, J. Zittartz: <i>A new method for the excitations of the one-dimensional Hubbard model</i> <i>Z. Phys. B</i> 78 (1990) 99.	75

-
- [9] M. Takahashi: *Magnetization curve for the half-filled Hubbard model*
Prog. Theor. Phys. **42** (1969) 1098. 86
- [10] K. Penc and F. Woynarovich: *Novel magnetic properties of the Hubbard chain with an attractive interaction*
Z. Phys. B **85** (1991) 269. 95
- [11] M. Takahashi: *One-Dimensional Hubbard model at finite temperature*
Prog. Theor. Phys. **47** (1972) 69. 107
- [12] M. Takahashi: *Low temperature specific heat of one-dimensional Hubbard model*
Prog. Theor. Phys. **52** (1974) 103. 121
- [13] F. H. L. Eßler, V. E. Korepin and K. Schoutens: *Completeness of the $SO(4)$ extended Bethe ansatz for the one-dimensional Hubbard model*
Nucl. Phys. B **384** (1992) 431. 133
- [14] B. S. Shastry: *Decorated star-triangle relations and exact integrability of the one-dimensional Hubbard model*
J. Stat. Phys. **50** (1988) 57. 161
- [15] T. Koma: *An extension of the thermal Bethe ansatz*
Prog. Theor. Phys. **83** (1990) 655. 184
- [16] H. Tsunetsugu: *Temperature dependence of spin correlation length of half-filled one-dimensional Hubbard model*
J. Phys. Soc. Japan **60** (1991) 1460. 190
- [17] A. M. Finkel'shtein: *Correlation functions in the one-dimensional Hubbard model*
Sov. Phys. JETP Lett. **25** (1977) 73. 194
- [18] H. J. Schulz: *Correlated fermions in one dimension*
Int. J. Mod. Phys. B **5** (1991) 57. 198
- [19] J. M. P. Carmelo and A. A. Ovchinnikov: *Generalization of the Landau liquid concept: example of the Luttinger liquid*
J. Phys. Cond. Mat. **3** (1991) 757. 216

-
- [20] F. Woynarovich and H. P. Eckle: *Finite-size corrections for the low-lying states of a half-filled Hubbard chain*
J. Phys. A **20** (1987) L443. 225
- [21] N. M. Bogoliubov and V. E. Korepin: *The role of quasi-one-dimensional structures in high- T_c superconductivity*
Int. J. Mod. Phys. B **3** (1989) 427. 232
- [22] F. Woynarovich: *Finite-size effects in a non-half-filled Hubbard chain*
J. Phys. A **22** (1989) 4243. 245
- [23] H. Frahm and V. E. Korepin: *Correlation functions of the one-dimensional Hubbard model in a magnetic field*
Phys. Rev. B **43** (1991) 5653. 259
- [24] B. S. Shastry and B. Sutherland: *Twisted boundary conditions and effective mass in Heisenberg-Ising and Hubbard rings*
Phys. Rev. Lett. **65** (1990) 243. 269
- [25] M. Fowler and N. Yu: *Persistent current of a Hubbard ring threaded with a magnetic flux*
Phys. Rev. B **45** (1992) 11795. 273
- II. The Supersymmetric t - J Model 283**
- [1] B. Sutherland: *Model for a multicomponent quantum system*
Phys. Rev. B **12** (1975) 3795. 287
- [2] P. Schlottmann: *Integrable narrow-band model with possible relevance to heavy-fermion systems*
Phys. Rev. B **36** (1987) 5177. 298
- [3] P. A. Bares and G. Blatter: *Supersymmetric t - J model in one dimension: separation of spin and charge*
Phys. Rev. Lett. **64** (1990) 2567. 307
- [4] S. Sarkar: *The supersymmetric t - J model in one dimension*
J. Phys. A **24** (1991) 1137. 311

[5]	N. Kawakami and S.-K. Yang: <i>Luttinger liquid properties of highly-correlated electron systems in one dimension</i> <i>J. Phys. C</i> 3 (1991) 5983.	326
[6]	A. Förster and M. Karowski: <i>Algebraic properties of the Bethe ansatz for an $sp(2,1)$-supersymmetric t-J model</i> <i>Nucl. Phys. B</i> 396 (1993) 611.	352
III.	Other Models of Strongly Correlated Electrons	381
A.	Models with Local Interaction	382
[1]	C. N. Yang: <i>Some exact results for the many-body problem in one dimension with repulsive delta-function interaction</i> <i>Phys. Rev. Lett.</i> 19 (1967) 1312.	384
[2]	M. Takahashi: <i>One-dimensional electron gas with delta-function interaction at finite temperature</i> <i>Prog. Theor. Phys.</i> 46 (1971) 1388.	388
[3]	D. C. Mattis, E. H. Lieb: <i>Exact solution of a many-fermion system and its associated boson field</i> <i>J. Math. Phys.</i> 6 (1965) 304.	407
[4]	F. D. M. Haldane: <i>'Luttinger liquid theory' of one-dimensional quantum fluids: I. Properties of the Luttinger model and their extension to the general 1D interacting spinless Fermi gas</i> <i>J. Phys. C</i> 14 (1981) 2585.	416
[5]	F. D. M. Haldane: <i>Demonstration of the "Luttinger Liquid" character of Bethe-ansatz-soluble models of 1-D quantum fluids</i> <i>Phys. Lett.</i> 81 A (1981) 153.	441
[6]	R. Z. Bariev: <i>Integrable spin chain with two- and three-particle interactions</i> <i>J. Phys. A</i> 24 (1991) L549.	444

-
- [7] R. Z. Bariev, A. Klümper, A. Schadschneider and J. Zittartz:
Exact solution of a one-dimensional model of hole superconductivity
J. Phys. A **26** (1993) 1249. 449
- [8] F. H. L. Eßler, V. E. Korepin and K. Schoutens:
New exactly solvable model of strongly correlated electrons
motivated by high T_c superconductivity
Phys. Rev. Lett. **68** (1992) 2960. 458
- B. Models with long-range interaction 462**
- [1] B. Sutherland: *Quantum many-body problem in one*
dimension: ground state
J. Math. Phys. **12** (1971) 246. 464
- [2] B. Sutherland: *Quantum many-body problem in one*
dimension: thermodynamics
J. Math. Phys. **12** (1971) 251. 469
- [3] Y. Kuramoto, H. Yokoyama: *Exactly soluble supersymmetric*
 t - J model with long-range exchange and transfer
Phys. Rev. Lett. **67** (1991) 1338. 475
- [4] N. Kawakami: *Asymptotic Bethe ansatz: Application to the*
one-dimensional t - J model with long-range exchange and transfer
Phys. Rev. B **45** (1992) 7525. 479
- References 483**