
CONTRIBUTORS

J.A. Andrades

*Department of Cell Biology, Faculty of Sciences
University of Málaga, Málaga, Spain*

J. Becerra

*Department of Cell Biology, Faculty of Sciences
University of Málaga, Málaga, Spain*

G. Benke

*Department of Orthopaedics
The Medical University of Warsaw, Warsaw, Poland*

J.P.A. van den Bergh

*St. Antonius Hospital
CM Nieuwegein, The Netherlands*

K. Bessho

*Department of Oral and Maxillofacial Surgery
Graduate School of Medicine, Kyoto University, Kyoto, Japan*

E. Birr

*Bone Transplantation Research Group
University of Oulu, Oulu, Finland*

T.R. Blattert

*Department of Trauma Surgery
Wuerzburg University Hospital, Wuerzburg, Germany*

C.M. ten Bruggenkate

*Rijnland Hospital
location St Elisabeth, CC Leiderdorp, The Netherlands*

S.K. Bulstra

*Department of Orthopaedic Surgery
Academic Hospital, AZ Maastricht, The Netherlands*

E.H. Burger

*Department of Oral Cell Biology
ACTA-Vrije Universiteit, Amsterdam, The Netherlands*

D. Chen

*Department of Medicine, Division of Endocrinology
University of Texas Health Science Center at San Antonio
San Antonio, Texas, USA*

F.L. Chen

*Department of Oral and Maxillofacial Surgery, Stomatological College
The Fourth Military Medical University
Xi'an, Shaanxi, China*

C. Chillemi

*Department of Orthopaedics
University of Ancona, Ancona, Italy*

S. Chubinskaya

*Department of Biochemistry
Rush Medical College at Rush Presbyterian–St.Luke's Medical Center
Chicago, IL, USA*

M.J. Danesh-Meyer

*Laboratory for Applied Periodontal and Craniofacial Regeneration
Department of Periodontology, Temple University School of Dentistry
Philadelphia, PA, USA*

G.H.G. Dekkers

*Department of Orthopaedic Surgery
Academic Hospital, AZ Maastricht, The Netherlands*

J.S. Erulkar

*Department of Orthopaedics and Rehabilitation
Yale University School of Medicine
New Haven, CT, USA*

C. Ferretti

*Bone Research Laboratory, MRC/University of Witwatersrand
Johannesburgh, South Africa*

T.J. Gao

*Department of Biomedical Engineering and Dentistry
Faculty of Medicine and Dentistry, University of Alberta
Edmonton, Alberta, Canada*

R.G.T. Geesink

*Department of Orthopaedic Surgery
Academic Hospital, AZ Maastricht, The Netherlands*

A. Gigante

*Department of Orthopaedics
University of Ancona, Ancona, Italy*

L.R. Giltaij

LV Doorn, The Netherlands

J.N. Grauer

*Department of Orthopaedics and Rehabilitation
Yale University School of Medicine
New Haven, CT, USA*

F. Greco

*Department of Orthopaedics
University of Ancona, Ancona, Italy*

L. Han

*Department of Oral and Maxillofacial Surgery, Stomatological College
The Fourth Military Medical University
Xi'an, Shaanxi, China*

O. Hanisch

*Department of Prosthodontics
University of Aachen Medical Center, Aachen, Germany*

I.C. Heyligers

*Department of Orthopaedic Surgery
University Hospital Vrije Universiteit
Amsterdam, The Netherlands*

O. Hietala

*Bone Transplantation Research Group
University of Oulu, Oulu, Finland*

N.H.M. Hoefnagels

*Department of Orthopaedic Surgery
Academic Hospital, AZ Maastricht, The Netherlands*

Y. Jin

*Department of Oral Pathology, Stomatological College
The Fourth Military Medical University
Xi'an, Shaanxi, China*

J. Klein-Nulend

*Department of Oral Cell Biology
ACTA-Vrije Universiteit, Amsterdam, The Netherlands*

K.E. Kuettner

*Department of Biochemistry
Rush Medical College at Rush Presbyterian–St. Luke's Medical Center
Chicago, IL, USA*

R. Kuijer

*Department of Orthopaedic Surgery
Academic Hospital, AZ Maastricht, The Netherlands*

K. Kusumoto

*Department of Plastic and Reconstructive Surgery
Kansai Medical University, Osaka, Japan*

M. Lewandowska-Szumiel

*Department of Transplantology and Central Tissue Bank
Center of Biostructure Research
Medical University of Warsaw, Warsaw, Poland*

X. Li

*Department of Oral Pathology, Stomatological College
The Fourth Military Medical Hospital
Xi'an, Shaanxi, China*

T.C. Lindholm

*Department of Oral and Maxillofacial Surgery
University Central Hospital of Turku, Finland*

E. Lourenco

*State University of Rio de Janeiro – UERJ
Rio de Janeiro, Brazil*

H. Lu

*Department of Oral Pathology, Stomatological College
The Fourth Military Medical Hospital
Xi'an, Shaanxi, China*

Q. Ma

*Department of Oral and Maxillofacial Surgery, Stomatological College
The Fourth Military Medical University
Xi'an, Shaanxi, China*

T.Q. Mao

*Department of Oral and Maxillofacial Surgery, Stomatological College
The Fourth Military Medical University
Xi'an, Shaanxi, China*

J.L. Ong

*University of Texas Health Science Center at San Antonio
Division of Biomaterials, San Antonio, Texas, USA*

M.M. Panjabi

*Department of Orthopaedics and Rehabilitation
Yale University School of Medicine
New Haven, CT, USA*

T.C. Patel

*Department of Orthopaedics and Rehabilitation
Yale University School of Medicine
New Haven, CT, USA*

A.H. Reddi

*Center for Regeneration and Repair
University of California, Davis, Sacramento
California, USA*

U. Ripamonti

*Bone Research Laboratory, MRC/University of the Witwatersrand
Johannesburg, South Africa*

H.F. Sailer

*Department for Cranio-Maxillofacial Surgery
University Hospital, Zurich, Switzerland*

A. Shimmin

*Melbourne Orthopaedic Group
Windsor Victoria, Australia*

X. Si

*Shanghai Research Institute of Stomatology
The Ninth Affiliated People's Hospital
Shanghai Second Medical University, Shanghai, China*

J.R. Tasker

*Bone Research Laboratory, MRC/University of Witwatersrand
Johannesburg, South Africa*

G.L. Tipoe

*Department of Anatomy, Faculty of Medicine
The University of Hong Kong, Hong Kong, China*

M. Ulmanen

*Bone Transplantation Research Group
University of Oulu, Oulu, Finland*

H. Uludag

*Department of Biomedical Engineering
Faculty of Medicine and Dentistry, University of Alberta
Edmonton, Alberta, Canada*

H.X. Wang

*Institute of Basic Medical Sciences
Academy of Military Medical Sciences, Beijing, China*

F.E. Weber

*Department for Cranio-Maxillofacial Surgery
University Hospital, Zurich, Switzerland*

U.M.E. Wikesjö

*Department of Periodontology
Temple University School of Dentistry
Philadelphia, PA, USA*

L. Yang

*Department of Oral Pathology, Stomatological College
The Fourth Military Medical University
Xi'an, Shaanxi, China*

S.L. Zhang

*Department of Oral and Maxillofacial Surgery, Stomatological College
The Fourth Military Medical University
Xi'an, Shaanxi, China*

M. Zhao

*Department of Medicine, Division of Endocrinology
University of Texas Health Science Center at San Antonio
San Antonio, Texas, USA*