

Contents

1. Introduction	1
2. Ueda's "Strange Attractors"	5
3. Pendulum	11
3.1. Equation of motion, linear and weakly nonlinear oscillations	11
3.2. Method of Poincaré map	18
3.3. Stable and unstable periodic solutions	20
3.4. Bifurcation diagrams	24
3.5. Basins of attraction of coexisting attractors	28
3.6. Global homoclinic bifurcation	33
3.7. Persistent chaotic motion — chaotic attractor	39
3.8. Cantor set — an example of a fractal geometric object	46
4. Vibrating System with Two Minima of Potential Energy	49
4.1. Physical and mathematical model of the system	50
4.2. The single potential well motion	53
4.3. Melnikov criterion	57
4.4. Fractal boundaries of basins of attraction and transient chaos in the region of principal resonance	62
4.5. Oscillating chaos and unpredictability of the final state after destruction of the resonant attractor	71
4.6. Boundary crisis of the oscillating chaotic attractor	75
4.7. Persistent cross-well chaos	79
4.8. Lyapunov exponents	82
4.9. Intermittent transition to chaos	84
4.10. Large Orbit and the boundary crisis of the cross-well chaotic attractor	87
4.11. Various types of attractors of the two-well potential system	94
5. Closing Remarks	98
Bibliography	101
Index	105