

Sums of independent random variables and Markov chains by Naresh Jain

Professor Kai Lai Chung's contributions to probability theory have had a major influence on several areas of research in the subject. I will restrict my comments to some of his work in two areas, sums of independent random variables and the theory of Markov chains, which led to a significant amount of further work, including some of my own.

Sums of Independent Random Variables

Kai Lai has made many outstanding contributions to this field, but I would like to concentrate on his 1948 paper [2]. If X_1, X_2, \dots is a sequence of real-valued independent random variables, and $S_n = X_1 + X_2 + \dots + X_n$, $n \geq 1$, denotes the sequence of partial sums, then the almost sure behavior of "large values" of $\{S_n\}$ was very well understood. Indeed, in the independent and identically distributed (i.i.d) case, Hartman and Wintner in 1941 [11] had already proved their celebrated law of the iterated logarithm: $EX_1 = 0$, $EX_1^2 = 1$, imply

$$\limsup_{n \rightarrow \infty} \frac{S_n}{(2n \log \log n)^{1/2}} = 1, \text{ a.s.} \quad (1)$$

In a more general non-i.i.d. context, Feller in 1943 [8] had written almost the final word. Chung [2] observed that if S_n is replaced by $|S_n|$ in (1), the assertion remains valid. However, if S_n^* denotes $\max_{1 \leq j \leq n} |S_j|$, then the behavior of "small values" of S_n^* had yet to be understood. He studied this problem in [2] including the non-i.i.d. situation and proved that if $EX_j \equiv 0$, and $E(S_n^2)$ is denoted by s_n^2 , then under a natural third moment assumption

$$\liminf_{n \rightarrow \infty} \frac{S_n^*}{s_n (\log \log s_n)^{-1/2}} = 8^{-1/2} \pi, \text{ a.s.} \quad (2)$$

To prove this result, he obtained the very profound probability estimate

$$P(S_n^* < cs_n) = \frac{4}{\pi} \sum_{j=0}^{\infty} \frac{(-1)^j}{2j+1} \exp\left(-\frac{(2j+1)^2 \pi^2}{8c^2}\right) + O((\log \log s_n / \log s_n)^{1/2}). \quad (3)$$

This contains the probability distribution for a standard one-dimensional Brownian motion process $\{B_t, t \geq 0\}$ if S_n^*/s_n is replaced by $\max_{0 \leq t \leq 1} |B_t|$ on the left-side and the second term is replaced by zero on the right-side.

In the i.i.d. case, two questions arose after Chung's work. The first one was raised by Chung himself: If $EX_1 = 0$, $EX_1^2 = 1$, does (2) hold without any further assumptions with $s_n^2 = n$? The second natural question was to obtain the analogue of (2) if X_1 is in the domain of attraction of a stable law.

As to the first question, several papers appeared on the subject getting close to the conditions stipulated by Chung. The question was finally settled in the affirmative by Jain and Pruitt in 1975 [16]. The probability distribution given in (3) by Chung played a key role in the final solution. For the second question, if the index of stability $\alpha < 2$, it was not even clear if one should expect an analogue of (2).

Fristedt had already observed that an analogue of (1) could not exist if $\alpha < 2$. However, Fristedt and Pruitt [9] and Jain and Pruitt [15] showed, under different conditions on X_1 , the existence of a real sequence $\{b_n\}$ increasing to infinity such that $\liminf(S_n^*/b_n) = c$, a.s., with $0 < c < \infty$, but it was not clear if the constant c depended on the distribution of X_1 or on the limit distribution alone.

Donsker and Varadhan [6] approached these problems through their large deviations probability estimates for stable processes and obtained explicit expressions for the limit constants. Jain [13] was then able to show through an invariance principle that the limit constant for the \liminf behavior of (S_n^*/b_n) is the same as for the relevant stable process obtained by Donsker and Varadhan [6].

The story by no means ends here. For a two-parameter Brownian motion $B(s, t)$, $0 \leq s, t \leq 1$, the leading term of the "small ball" probability estimate for $P(\max_{0 \leq s, t \leq 1} |B(s, t)| \leq c)$, as $c \downarrow 0$, is of great interest and turned out to be a challenging problem, which was solved by Bass [1] and Talagrand [21]. Much work has been done by other authors for parameter dimension larger than 2; these results, however, are not as definitive as in the two-dimensional parameter case. Many difficult questions still remain to be answered and we can expect these investigations to continue, owing their origins to Chung's pioneering work.

Markov Chains

It is difficult to imagine that anybody working in the area of Markov processes would not be familiar with Chung's monograph: Markov Chains with Stationary Transition Probabilities [3]. This monograph deals with countable state Markov chains in both discrete time (Part I) and continuous time (Part II). Much of Kai Lai's fundamental work in the field is included in this monograph. My comments will be confined to Part I. Here, for the first time, Kai Lai gave a systematic exposition of the subject which includes classification of states, ratio ergodic theorems, and limit theorems for functionals of the chain.

For a general state space, Doeblin had given a classification scheme in a seminal paper in 1937 [5]. This and other work of Doeblin had a major impact on the field and led to further developments by Chung [4], Doob [7], Harris [10], and Orey [18], [19]. In the early sixties there were a number of basic ingredients of a general state space theory that would lead to an exact counterpart of Part I of Chung [3]. Much fundamental ground work, including positive recurrence (the so-called Doeblin's condition), was done by Doob [7]. Harris [10] introduced his recurrence condition: There exists a nonzero σ -finite measure φ on the state space S such that $\varphi(E) > 0$ implies that starting from every $x \in S$, E is visited infinitely often a.s. He proved the existence of a (unique) σ -finite invariant measure π under this condition. If $\pi(S) = +\infty$, one could call the process null-recurrent, and one could ask if $\pi(E) < \infty$ implied $P^n(x, E) \rightarrow 0$ for every $x \in S$, as $n \rightarrow \infty$; here $P^n(x, E)$ denotes the n -step transition probability from x to E . This result was conjectured by Orey [18] and was a natural extension to the general state space situation of the corresponding well-known result for a countable state chain. Under Harris's recurrence condition one could also ask if an analogous ratio ergodic theorem was true, namely, if $\pi(F) > 0$ and $\pi(E) < \infty$, does

$$\sum_{j=1}^n P^j(x, E) / \sum_{j=1}^n P^j(y, F) \rightarrow \pi(E) / \pi(F) \quad (4)$$

as $n \rightarrow \infty$ for all $x, y \in S$? These questions were answered in [12] under Chung's guidance. Chung gave an example, reported in [12], to show that in the general case, as opposed to the countable state space case, (4) is true only for π -almost all x, y , and not for all x, y . The situation is different when the state space is not countable because one could stay in a π -null set for a rather long time! A little later, Jain and Jamison [14] introduced an irreducibility condition: There exists a nonzero σ -finite measure φ on S such that $\varphi(E) > 0$ implies that starting from every $x \in S$, the process visits E with positive probability. In this paper they essentially brought the program of Doeblin [5] and Chung [4] to completion. Chung's influence can be seen throughout these works. For other work in the area one can refer to monographs by Orey [19], Revuz [20] and Meyn and Tweedie [17].

References

- [1] Bass, R. (1988). Probability estimates for multiparameter Brownian processes, *Ann. Probab.* **16**, 251-264.
- [2] Chung, K.L. (1948). On the maximum partial sums of independent random variables, *Trans. Amer. Math. Soc.* **64**, 205-233.
- [3] Chung, K.L. (1960). *Markov chains with stationary transition probabilities*, *Grund. Math. Wiss.* **104**, Springer-Verlag, New York/Berlin. Second Edition (1967).

- [4] Chung, K.L. (1964). The general theory of Markov processes according to Doebelin, *Z. Wahr. Verw. Gebiete* **2**, 230-254.
- [5] Doebelin, W. (1937). Sur les propriétés asymptotiques de mouvement régis par certain type de chaines simples. *Bull. Math. Soc. Roum. Sci.* **39**, No. 1, 57-115; No. 2, 3-61.
- [6] Donsker, M.D. and Varadhan, S.R.S. (1977). On laws of the iterated logarithm for local times, *Comm. Pure Appl. Math.* **30**, 707-753.
- [7] Doob, J.L. (1953). *Stochastic Processes*, Wiley and Sons, New York.
- [8] Feller, W. (1943). The general form of the so-called law of the iterated logarithm, *Trans. Amer. Math. Soc.* **54**, 373-402.
- [9] Fristedt, F. and Pruitt W.E. (1971). Lower functions for increasing random walks and subordinators, *Z. Wahr. Verw. Gebiete* **18**, 167-182.
- [10] Harris, T.E. (1955). The existence of stationary measures for certain Markov processes, *Proc. Third Berkeley Symp. Math. Statist. Probab. Vol II*, 113-124.
- [11] Hartman, P. and Wintner, A. (1941). On the law of the iterated logarithm, *Amer. J. Math.* **63**, 169-176.
- [12] Jain, N.C. (1966). Some limit theorems for a general Markov process, *Z. Wahr. Verw. Gebiete* **6**, 206-223.
- [13] Jain, N.C. (1982). A Donsker-Varadhan type of invariance principle, *Z. Wahr. Verw. Gebiete* **59**, 117-138.
- [14] Jain, N.C. and Jamison, B. (1967). Contributions to Doebelin's theory of Markov processes, *Z. Wahr. Verw. Gebiete* **8**, 19-40.
- [15] Jain, N.C. and Pruitt, W.E. (1973). Maxima of partial sums of independent random variables, *Z. Wahr. Verw. Gebiete* **27**, 141-151.
- [16] Jain, N.C. and Pruitt, W.E. (1975). The other law of the iterated logarithm, *Ann. Probab.* **3**, 1046-1049.
- [17] Meyn, S.P. and Tweedie, R.L. (1993). *Markov chains and stochastic stability*, Springer-Verlag, New York.
- [18] Orey, S. (1959). Recurrent Markov chains, *Pacific J. Math.* **9**, 805-827.
- [19] Orey, S. (1971). *Limit Theorems for Markov Chain Transition Probabilities*, Van Nostrand, New York.
- [20] Revuz, D. (1975). *Markov Chains*, North-Holland, New York. Revised Edition (1984).
- [21] Talagrand, M. (1994). The small ball problem for the Brownian sheet, *Ann. Probab.* **22**, 1331-1354.

Naresh Jain
School of Mathematics
University of Minnesota